

PROSE

MAPP 19196

Contains 800 g/L prosulfocarb as an emulsifiable concentrate formulation. For the control of annual grass weeds and annual broad-leaved weeds in Winter Barley, Winter Wheat and Maincrop potatoes. The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

PROTECT FROM FROST
SHAKE WELL BEFORE USE

PROSE
UFI Code: XXXX



Danger

May be fatal if swallowed and enters airways

Causes skin irritation

May cause an allergic skin reaction

Causes serious eye irritation

Very toxic to aquatic life with long lasting effects

Avoid breathing dust/fume/gas/mist/vapours/spray.

Wear protective gloves/protective clothing/eye protection/face protection

IF SWALLOWED: immediately call a POISON CENTER or doctor/physician

Do NOT induce vomiting.

If skin irritation or rash occurs: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste

To avoid risks to human health and the environment, comply with the instructions for use

IMPORTANT INFORMATION
FOR USE ONLY AS A PROFESSIONAL AGRICULTURAL HERBICIDE

Crops	Maximum individual dose (L product/Ha)	Maximum number of treatments (per crop)	Latest time of application
Wheat (winter), barley (winter)	5 OR	1	Pre-crop emergence
	3	1	Early tillering (GS21)
Potato	5	1	At emergence (soil rising over emerging potato shoots)

Other specific restrictions:

Do not apply by hand-held equipment

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate or handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before eating and after work.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with the 'Local Environment Risk Assessment for Pesticides Horizontal Boom Sprayers' booklet available from the HSE Chemicals Regulation Division's website or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Extreme care must be taken to avoid spray drift onto non-crop plants outside of the target area.

Contents: 5 – 20 L
Batch number: See container

Authorisation holder
Sharda Agrochem Limited
201 Cervantes House
5-9 Headstone Road
Harrow, Middlesex
HA1 1PD, United Kingdom



Sharda Agrochem LTD

Marketing company
Sharda Cropchem España S.L.
Edificio Atalayas Business Center
Carril Condomina nº 3, Planta 12
30006 Murcia, Spain

24-HOUR EMERGENCY TELEPHONE NUMBER: +44 (0) 1865 407333

DIRECTIONS FOR USE

IMPORTANT: this information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

RESTRICTIONS

Do not apply to crops under stress or to crops suffering from waterlogging, pest attack, disease, frost or the effects of high diurnal temperature changes. Transient yellowing can occur although crops fully recover.
For cereals, seed must be covered by 3cm of soil and for best results apply to a firm, moist seedbed free of clods.
DO NOT sow field beans or broad beans within 12 months of application.

WEEDS CONTROLLED

GRASSWEEDS CONTROLLED	
Susceptible	
Rough Stalked Meadow grass	Pre-emergence
Moderately susceptible	
Annual meadow grass	Up to 3 true leaves
Loose silky bent	Pre-emergence

PROSE used pre-emergence will reduce blackgrass and Italian ryegrass populations. It should only be used as part of an appropriate management strategy involving sequences with products of alternative modes of action and the use of cultural techniques.

BROAD-LEAVED WEEDS CONTROLLED	
Susceptible	
Ivy leaved speedwell	Susceptible at cotyledon stage
Black nightshade	Pre-emergence
Chickweed	Susceptible at up to 2 true leaves
Forget me not	Pre-emergence
Cranesbill	Pre-emergence
Red dead nettle	Pre-emergence
Common field speedwell	Susceptible at cotyledon stage
Green field speedwell	Susceptible at cotyledon stage
Wall speedwell	Pre-emergence

WEED RESISTANCE

Strains of some annual grasses (e.g. black-grass, wild-oats, and Italian rye-grass) have developed resistance to a range of herbicides which may lead to poor control from one or more product or mode of action. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop adviser or product manufacturer. PROSE should only be used for control of herbicide resistant strains of annual grasses as part of an appropriate management strategy, including sequences with herbicides of alternative modes of action and the use of cultural techniques.

CROP SPECIFIC INFORMATION

Winter Cereals

PROSE applied pre-emergence may infrequently slow crop emergence. This effect is transient and has been demonstrated not to adversely affect yield.

Potatoes

For control of a wider spectrum of weeds use PROSE in mixture with either RETRO or approved formulations of products containing the active ingredient metribuzin, only as a pre-emergence application. Please consult company literature for specific product and best use guidelines.
Always observe full label restrictions for any tank mix partner.

TIMING

Winter Cereals

PROSE can either be used at pre-emergence of the crop (5 L/ha) or at post-emergence of the crop up to GS 21 (3 L/ha). One application per crop.

Early and Maincrop Potatoes

PROSE may be applied pre-emergence or at emergence of the crop (soil rising over emerging potato shoots) at 5L/ha. Complete ridge formation before application of PROSE and do not disturb treated soil after application.

FOLLOWING CROPS

Winter Cereals, Early and Maincrop Potatoes

Do not sow field beans or broad beans within 12 months of application.

In the case of winter cereal crop failure, Winter Wheat or Winter Barley may be sown immediately in the autumn.

The following crops may be sown in the spring after winter cereal crop failure:

- **Without ploughing:** Sunflowers, Maize, Flax, Spring oats, Spring barley, Spring wheat, Spring peas, Spring oilseed rape, Soya beans.
- **With ploughing:** Carrots, Lettuce, Onions, Sugar beets, Potatoes.
- **Do not sow:** Field beans, Broad beans.

The following crops may be sown in the autumn after potato crop failure or normal harvest:

Without ploughing: Winter oats, Winter barley, Winter wheat

MIXING

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of PROSE and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying.

SPRAYING

Apply PROSE in a water volume of 200-400 litres per hectare.

Apply PROSE using a conventional fan nozzle producing a medium spray quality as Defined by the British Crop Protection Council. A spray pressure of 2.0-3.0 bars is recommended.

PROSE is rainfast after 1 hour.

WASHING OUT PROCEDURE

Immediately after use, clean the spray equipment thoroughly. Drain the system completely and rinse spray tank, boom and nozzles two to three times with clean water until the foam and all traces of product have been removed.

COMPANY ADVISORY INFORMATION

The following does not form part of the product label under Regulation (EC) No 1107/2009 and provides additional advice on the product use at the discretion of Sharda.

Apply with a Droplet Spectrum of Coarse to Very Coarse, use 3* LERAP (minimum 75% Drift Reduction Nozzles) at correct drift reduction operating pressures.

Keep forward speeds < 12 km/hr.

Ensure Boom height is maintained @ 50cm above target.

Apply at 200 l water/ha.

Wind < 4 m/s.

Do not apply in still wind conditions (Force 0)

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable but, as we cannot exercise control over their storage, handling, mixing or use or the weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us or re-sellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.